

Patient: [REDACTED] Specimen ID: [REDACTED]
 DOB: [REDACTED] Patient ID: D [REDACTED] Control ID: [REDACTED] Date collected: [REDACTED]

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
T4, Free (Direct)	1.21		ng/dL	0.82 - 1.77	01
DHEA-Sulfate	141.4		ug/dL	71.6 - 375.4	01
TSH	2.090		uIU/mL	0.450 - 4.500	01
Prolactin	6.3		ng/mL	4.0 - 15.2	01
Prostate-Specific Ag, Serum					
Prostate Specific Ag, Serum	1.3		ng/mL	0.0 - 4.0	01
Roche ECLIA methodology. According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.2 ng/mL or greater followed by a subsequent confirmatory PSA value 0.2 ng/mL or greater. Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.					
IGF-1					
Insulin-Like Growth Factor I	151		ng/mL	61 - 200	02
Estradiol, Sensitive	35.3	High	pg/mL	8.0 - 35.0	02
This test was developed and its performance characteristics determined by LabCorp. It has not been cleared by the Food and Drug Administration. Methodology: Liquid chromatography tandem mass spectrometry (LC/MS/MS)					
Triiodothyronine (T3), Free	2.8		pg/mL	2.0 - 4.4	01
Sex Horm Binding Glob, Serum	38.1		nmol/L	19.3 - 76.4	01

Comments:

^A This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

01	MB	LabCorp Birmingham 1801 First Avenue South, Birmingham, AL 35233-1935	Dir: Brian Ragland, MD
02	BN	LabCorp Burlington 1447 York Court, Burlington, NC 27215-3361	Dir: Sanjai Nagendra, MD
03	ES	Esoterix Inc 4301 Lost Hills Road, Calabasas Hills, CA 91301-5358	Dir: Samuel Pepkowitz, MD

For inquiries, the physician may contact Branch: 800-877-5227 Lab: 205-581-3500